



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

stretch from surface to surface without intercellular spaces. Consequently the width of the lines of starch produced at the edge of the darkened region is not widened even though the unlighted area is supplied with CO_2 under abnormal pressure. If the net be coarse the zone of marginal starch will be wider than if it is fine. Diffusion of self-produced CO_2 to 2.5 cm at most is thus possible; for in parallel-veined leaves of *Triticum*, *Hordeum*, and *Zea*, though the veins do not prevent diffusion, the intercellular passages are so narrow as to limit it to 3 cm . In *Tradescantia* and *Acorus* the transverse anastomoses prevent more extensive movement. In *Eichhornia*, *Pontederia*, and *Eucomis* the leaves have spacious intercellular passages, and so the movement is much more free. But even here the diffusion scarcely surpasses 3 cm , unless through a region of the leaf that is not in condition to act on the CO_2 . In nature, therefore, movement of CO_2 may be considered practically nil.—C. R. B.

Seedling of a graft-hybrid.—Certain branches of the graft hybrid, *Cytisus Adami*, revert, producing flowers having the characters of the reputed parents *C. Laburnum* and *C. purpureus* respectively. The *C. Adami* flowers are ordinarily sterile, while those borne on reverted branches reproduce their respective parents.

In May, 1904, HILDEBRAND³⁵ observed that several flowers of a *C. Adami* branch of a cultivated specimen in the Freiburg botanical garden had set seed, and was able to obtain three fruits from them, which had chiefly the characters of *C. Laburnum*, but in certain respects resembled *C. purpureus*. It is not known whether these flowers were self-pollinated, but it is not unlikely that the pollen came from *C. Laburnum* flowers, since the *C. Adami* flowers are usually sterile. Two of the seeds germinated. Both were very similar in character to *C. Laburnum* and in 1907 one of them produced hundreds of flowers, all having the characters of *C. Laburnum*. No conclusions can be drawn regarding the hereditary bearing of these facts, in the absence of a knowledge of the manner of pollination of the flowers and the nature of the next generation of offspring.—R. R. GATES.

Chlorophyll.—The discussion as to the phosphorus content of chlorophyll waxes warm. STOKLASA replies vigorously³⁶ to TSWETT's criticisms³⁷ and takes issue with WILLSTÄTTER's results.³⁸ The question is yet in the stage of polemic

³⁵ HILDEBRAND, FRIEDRICH, Ueber Sämlinge von *Cytisus Adami*. Ber. Deutsch. Bot. Gesells. **26a**:590-595. 1908.

³⁶ STOKLASA, J., BRALIK, V., UND ERNST, A., Zur Frage des Phosphorgehaltes des Chlorophylls. Ber. Deutsch. Bot. Gesells. **27**:10-20. 1909.

³⁷ TSWETT, M., Ist der Phosphor an dem Anbau des Chlorophylline beteiligt? *Ibid.* **26a**:214-220. 1908.

³⁸ WILLSTÄTTER, R., Zur Kenntniss der Zusammensetzung des Chlorophylls. Liebig's Annalen der Chemie **350**:48-82. 1906.

WILLSTÄTTER, R., UND BENZ, M., Ueber krystallisiertes Chlorophyll. *Ibid.* **358**:267-287. 1907.